



Entrez PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed



for

Go

Clear

☒ Limits

Preview/Index

History

Clipboard

Details

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Field: Title, Limits: Publication Date from 1980 to 2001

Display

Abstract



Show: 20



Sort



Send to

Text

☐ 1: Nat Med. 1996 May;2(5):545-50.

Related Articles, Links

## Immune responses to transgene-encoded proteins limit the stability of gene expression after injection of replication-defective adenovirus vectors.

Tripathy SK, Black HB, Goldwasser E, Leiden JM.

Department of Pathology, University of Chicago, Illinois 60637, USA.

The use of replication-defective adenoviruses (RDAd) for human gene therapy has been limited by host immune responses that result in transient recombinant gene expression in vivo. It remained unclear whether these immune responses were directed predominantly against viral proteins or, alternatively, against foreign transgene-encoded proteins. In this report, we have compared the stability of recombinant gene expression in adult immunocompetent mice following intramuscular (i.m.) injection with identical RDAd encoding self (murine) or foreign (human) erythropoietin. Our results demonstrate that immune responses direct against foreign transgene-encoded proteins are the major determinants of the stability of gene expression following i.m. injection of RDAd. Moreover, we demonstrate long-term recombinant gene expression in immunocompetent animals following a single i.m. injection of RDAd encoding a self protein. These findings are important for the design of future preclinical and clinical gene therapy trials.

PMID: 8616713 [PubMed - indexed for MEDLINE]

Display

Abstract



Show: 20



Sort



Send to

Text

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Jan 12 2005 06:52:28

Br Med Bull. 1995 Jan;51(1):205-16.  
Gene therapy and viral vaccination: the interface.  
Wilkinson GW, Borysiewicz LK.

Live viral vaccines have had a major impact on the incidence of acute virus infections world-wide. Virus infections recognised as future vaccine targets will require a modified approach based on a detailed understanding of the immunobiology of specific infections combined with the application of new technologies designed to generate specific and appropriate protective immunity. A similar vector technology directed at in vivo gene delivery is currently being exploited both for gene therapy and vaccination. The induction of an immune response to an expressed transgene represents a potential hazard for a gene therapy protocol but is the object of a vaccine strategy. In vivo gene delivery using replication-competent or replication-deficient viral vector systems and by direct transfer of naked DNA can generate an effective humoral, secretory and cell-mediated immune response to expressed transgenes.